

# **Viral Hepatitis Transmission in the United States**

---

**Ian T. Williams, PhD, MS**

Division of Viral Hepatitis

Centers for Disease Control and Prevention

Atlanta, GA

The findings and conclusions in this presentation have not been formally disseminated by CDC and should not be construed to represent any agency determination or policy




# Features of HBV & HCV Infection

		<u>HCV</u> <u>Infection</u>	<u>HBV</u> <u>Infection</u>
Pathogen		RNA virus	DNA virus
Incubation period	Average:	6–7 wks	8–12 wks
	Range:	2–26 wks	6–26 wks
Clinical illness (jaundice)		20%–30%	< 5 yrs: <10% > 5 yrs: 30–50%
Chronic hepatitis		~70%	< 5 yrs: 30–90% Older children: 6–10% Adults: <5%

# Disease Burden from Hepatitis B & Hepatitis C in the United States

<u>Outcome</u>	<u>HBV</u>	<u>HCV</u>
Percent ever infected	4.9%	1.6%
# chronic infections	~1.2 million	~3.2 million
# new infections/yr	~60,000	~30,000
# deaths/yr	~5,000	~10,000

# Acute Viral Hepatitis, United States, 2001-2004



Source: Sentinel Counties Study, CDC



# Transmission of Bloodborne Viral Infections

## Route

- **Percutaneous**

- Apparent

- Inapparent

- **Permucosal**

## Mode

injection drug use  
needle stick injury

blood/ serous fluid

sex  
perinatal

# Relative Transmission Efficiency of Bloodborne Viral Infections


	<b>HBV</b>	<b>HCV</b>	<b>HIV</b>
<b>Injection drug use</b>	+++	++++	++
<b>Sexual</b>	+++	+	++
<b>Perinatal</b>	++++	+	++
<b>Occupational</b>	+++	+/-	+/-

---

# **Epidemiology of Acute Hepatitis B in the United States**

---


# Estimated Number of Acute HBV Infections, United States, 1980-2004





# Reported Risk Factors for Acute Hepatitis B, United States, 2001-2004

---



<sup>1</sup> Other: Household contact, institutionalization, hemodialysis, blood transfusion, occupational exposure

Source: Sentinel Counties Study of Acute Viral Hepatitis, CDC

# Transfusion Associated Hepatitis B Cases Reported in the National Notifiable Disease Surveillance System, 2003

---

- Validation study of cases of acute symptomatic hepatitis B reported to the CDC from State health departments with transfusion reported as a risk factor for infection
- Of 7526 reported cases, 49 were reported with “transfusion” as a risk factor for infection
  - Only 10 had acute hepatitis B and were transfused during the exposure period
  - Only 1 with an infected donor (who was in the “window period” of infection)

---

# **Recent Case Reports of HBV Infection Following Blood Transfusion**

---

# Case Report 1: HBV Infection Following Blood Transfusion: New York, 2004

- A 60 year old woman developed acute hepatitis B in Sept, 2004 and died
  - No traditional risk factors for infection
  - Received four units packed red blood cells in May, 2004
- All four donors tested
  - One donor found to have become infected with HBV since donation
    - He admitted multiple male sex partners in 3 months prior to donation which was not disclosed at donation
    - No archived specimen for testing
- Likely donor was in the early incubation period of HBV infection

# Case Report 2: HBV Infection Following Blood Transfusion: Texas, 2004

- Repeat donor was found to be HBsAg positive in August, 2004
  - Previous donation in June, 2004 (HBsAg & anti-HBc negative) was traced to single recipient
- Recipient developed acute hepatitis B in Sept, 2004
- Donor did not disclose any risk factors at time of donation in June or August or upon re-interview
- Likely donor was in the early incubation period of HBV infection

# Implications of Recent Investigations of HBV Infection Following Blood Transfusion


- Transfusion transmitted HBV infection is a rare event
  - Risk of collecting HBV infectious blood in the window period is 1 in ~200,000 donations
  - When transmission has been observed, it has been due to “window period” donations and not testing errors
- Donor deferrals based on geographic, medical and behavioral factors are a first line of defense
  - Depends on donor honesty

# Immunization Strategy to Eliminate Transmission of HBV Infection in the United States

---

- Universal vaccination of all infants beginning at birth
- Prevention of perinatal HBV infection through
  - Routine screening of all pregnant women for HBsAg, and
  - Immunoprophylaxis of infants born to HBsAg positive women and infants born to women with unknown HBsAg status
- Routine vaccination previously unvaccinated children and adolescents
- Vaccination of previously unvaccinated adults at increased risk of infection

# Hepatitis B Vaccine 3 Dose Coverage Among 19-35 Month Old Children, by Year of Survey, 1992-2004\*




\*Source: National Immunization Survey





# Adult Hepatitis B Vaccine Coverage, 2002



National Health Interview Survey



# Estimated Hepatitis B Vaccine Coverage in Adults

---

Risk Group	Coverage Estimate (year, site)
Dialysis patients	60% (2001, national)
Occupationally-exposed workers	75% (2002-3, national)
Men who have sex with men	32% (1998-2000, YMS*)
Injection drug users	40% (2002-2004, DUIT**)
STD clinic clients	10% (1998-2001, San Diego)

\* Young Men's Survey: participants 22-29 years of age

\*\* Drug User Intervention Trial: participants 15-30 years of age




---


# **Epidemiology of Acute Hepatitis C in the United States**

---

# Incidence of Acute HCV Infection, United States



# Reported Risk Factors for Acute Hepatitis C, United States, 2001-2004



•\*Aggregate risk factor = Case shown a list of risk factors and admits to one, but does not specify which one

Source: Sentinel Counties Study of Acute Viral Hepatitis, CDC




# Transfusion Associated Hepatitis C Cases Reported in the National Notifiable Disease Surveillance System, 2003

---


- Validation study of cases of acute symptomatic hepatitis C reported to the CDC from State health departments with transfusion reported as a risk factor for infection
- Of 891 reported cases, 16 were reported with “transfusion” as a risk factor for infection
  - Only 1 had acute hepatitis C and was transfused during the exposure period
    - Received blood/blood products from six donors
    - Four of six donors re-tested and found to be uninfected

# Posttransfusion Hepatitis, United States



Alter HJ and Houghton M. Hepatitis C virus and eliminating post-transfusion hepatitis. *Nature Medicine* 2000;6:1082-6.

# Prevalence of HCV & HBV Infection Among Injection Drug Users in Two Time Periods



Garfein RS *Am J Public Health* 1996; 86:655 & Collaborative Injection Drug User Study III (CIDUSIII)/Drug Users Intervention Trial (DUI): Baltimore, Chicago, Los Angeles, New York City, Seattle. CDC unpublished data.





---

# **Recent Reports of Clusters of Acute HCV Infection Among Men who have Sex with Men (MSM) in Europe**

---

# Sexual Transmission of HCV

---

- Sexual transmission of HCV occurs, but efficiency is low
  - Rare between long-term steady partners
  - MSM at no higher risk than sexually active heterosexuals
- Factors that facilitate transmission between partners unknown (e.g., viral titer, other STDs)

# Clusters of Acute HCV Infection Among MSM in Europe

- Case report 1: France
  - Five HIV-infected MSM with acute HCV infection identified at a single clinic in a 13 month period
    - Denied IDU or other parenteral risk factors for infection
    - All reported unprotected anal intercourse and had concomitant syphilis
- Case report 2: The Netherlands
  - Seven MSM with acute HCV infection identified among 15 sexual contacts
    - Denied IDU or other parenteral risk factors for infection
    - All reported unprotected anal intercourse and sexual practices that included fisting
    - 6 of 7 cases had concomitant rectal lymphogranuloma venereum (LGV)
    - 6 of 7 cases were HIV infected

# Clusters of Acute HCV Infection Among MSM in Europe (II)

---

- No such clusters identified in the United States
  - Rare event or under reporting?
  - Difficult to study
  - Role of unreported IDU
- Need for further studies

---

# **Estimates of Past HBV and HCV Infection in Selected Populations in the United States**

---


# Caveats About Estimates of HBV and HCV Infection in Selected Populations

- Incidence does not equate with prevalence
  - Age/cohort/period effects: Changes in disease rate according to age, year of birth, & point in calendar time
  - In-migration and out-migration
- Geographic differences in prevalence
  - Urban v. rural or city v. city
- Individuals may have multiple risk factors
  - Belong to several population subgroups (e.g. MSM and IDU)
- External Validity: Ability to generalize results to “larger” population of interest
  - Where/how subjects selected (STD clinic v. street recruit)

# More Caveats About Estimates of HBV and HCV Infection in Selected Populations

- Prevalence of past infection v. prevalence of chronic infection
  - ~10% of persons in the US with past HBV infection (anti-HBc+) have chronic infection (HBsAg+)
    - In the general US population:
      - 4.9% of persons have evidence of past infection (anti-HBc+)
      - 0.4% have evidence of chronic infection (HBsAg+)
  - ~75% of persons with past HCV infection (anti-HCV+) have chronic infection (HCV RNA+)
    - In the general US population:
      - 1.6% of persons have evidence of past infection (anti-HCV+)
      - 1.3% have evidence of chronic infection (HCV RNA+)

# Estimates of Past HBV (anti-HBc) & HCV (anti-HCV) Infection in Selected US Populations





# Summary

- Incidence of acute HBV and HCV infection has declined in the past two decades
  - Primary risk factors remain unchanged
  - Impact of hepatitis B vaccination
- Transfusion historically was an important risk factor for infection (especially for HCV infection), but currently is extremely rare
  - When transmission has been observed, it has been due to “window period” donations and not testing errors
- Prevalent infections more common than acute infections
- Prevalence lower in younger age groups

# **CDC's Division of Viral Hepatitis**

---

**<http://www.cdc.gov/hepatitis>**